

Very Rare Flucloxacillin-Paracetamol Interaction risk of Metabolic Acidosis



Quick Read

Especially in patients with serious renal impairment, malnutrition or sepsis, concomitant use of flucloxacillin and maximum doses of paracetamol may very rarely be associated with high anion gap metabolic acidosis (HAGMA).

Flucloxacillin is a semi-synthetic isoxazolilpenicillin. This antibiotic has been used in Portugal since the 1970s to treat infections mainly caused by *Streptococci* and *Staphylococci*. Through various pathways, both flucloxacillin and paracetamol can cumulatively lead to an increase in 5-oxoproline (also known as pyroglutamic acid, pidolic acid or pyroglutamate).

Following an assessment of literature,¹⁻⁷ EudraVigilance (European ADR report database) and MA holder data, the European Medicines Agency (EMA) has confirmed an association between a rare type of metabolic acidosis and concomitant use of flucloxacillin and paracetamol: **high anion gap metabolic acidosis (HAGMA)**, which results from accumulation of pyroglutamic acid (5-oxoproline). Flucloxacillin inhibits 5-oxoprolinase activity resulting in the accumulation of 5-oxoproline. Paracetamol on the other hand, when used **chronically** or in **high** doses, can cause glutathione depletion, which in turn activates γ -glutamylcysteine synthetase, producing more γ -glutamylcysteine and consequently more circulating 5-oxoproline. Clearance of the latter is decreased in renal failure, which aggravates the condition. Other risk factors may also interfere with this metabolic cycle.¹

Though very rare, this condition is serious and potentially fatal. It is all the more relevant because paracetamol is available for self-medication as over-the-counter products.

In order to inform healthcare professionals about this risk, EMA has deemed necessary that a warning be included in the SmPC to underscore those **groups of patients who are at increased risk** of developing the condition (the PIL will be updated accordingly):

4.4. Special warnings and precautions for use

Caution is advised when flucloxacillin is administered concomitantly with paracetamol due to the increased risk of high anion gap metabolic acidosis (HAGMA). Patients at high risk for HAGMA are in particular those with severe renal impairment, sepsis or malnutrition especially if the maximum daily doses of paracetamol are used. After co-administration of flucloxacillin and paracetamol, a close monitoring is recommended in order to detect the appearance of acid-base disorders, namely HAGMA, including the search of urinary 5-oxoproline. If flucloxacillin is continued after cessation of paracetamol, it is advisable to ensure that there are no signals of HAGMA, as there is a possibility of flucloxacillin maintaining the clinical picture of HAGMA (see section 4.5).

4.5. Interaction with other medicinal products and other forms of interaction

Caution should be taken when flucloxacillin is used concomitantly with paracetamol as concurrent intake has been associated with high anion gap metabolic acidosis, especially in patients with risk factors. (see section 4.4.).

4.8. Undesirable effects

[...] Post marketing experience: very rare cases of high anion gap metabolic acidosis, when flucloxacillin is used concomitantly with paracetamol, generally in the presence of risk factors (see section 4.4.)

References:

- ¹ Hunter RW et al. Pyroglutamic acidosis in association with therapeutic paracetamol use. *Clinical Medicine*. 2016 Vol 16. No 6:524-9
- ² Jessurun N et al. Advanced age and female sex as risk factors for high anion gap metabolic acidosis after a drug interaction between paracetamol and flucloxacillin: a case series. *J Am Geriatr Soc*. 2016 Oct; 64(10):e90-e93.
- ³ Arnold JFH et al. Een metabole acidose met een verhoogde anion-gap ten gevolge van een zeldzame complicatie van de behandeling van een *Staphylococcus aureus*-bacteriëmie. *Tijdschr Infect* 2013;8 (6): 202-208.
- ⁴ Lanoy C, Bouckaert Y. Metabolic acidosis and 5-oxoprolineuria induced by flucloxacillin and acetaminophen: a case report. *J Med Case Rep* 2016 Jun 23;10(1):184.
- ⁵ Demsey et al. Pyroglutamic acidemia: A cause of high anion gap metabolic acidosis. *Crit Care Med* 2000; 28: 1803-1807.
- ⁶ Pitt JJ, Hauser S. Transient 5-oxoprolineuria and high anion-gap metabolic acidosis: clinical and biochemical findings in eleven subjects. *Clin Chem* 1998;44:1497-1503.
- ⁷ Van den Bergh FA et al. 5-Oxoprolineurie: een verworven stofwisselingsstoornis door behandeling met flucloxacilline en paracetamol. *Ned Tijdschr Klin Chem Labgeneesk* 2008; 33:39-42.

Ana Sofia Martins

INDEX CARD

Director: Fátima Canedo

Editor: Rui Pombal

Contributors: Ana Sofia Martins, António Leandro Ponte, Cristina Mousinho, Elsa de Fátima Costa, Fátima Bragança, Fátima Hergy, Leonor Nogueira Guerra, Magda Pedro, Márcia Silva, Sílvia Duarte, Vanda Araújo

Publishing Assistant: Inocência Pinto

Advisory Board: Conselho Diretivo do INFARMED, I.P. – Comissão de Avaliação de Medicamentos
INFARMED – Autoridade Nacional do Medicamento e Produtos de Saúde, I.P.
Parque de Saúde de Lisboa, Av. do Brasil, N.º 53, 1749-004 Lisboa

Phone: +351 217 987 100

E-mail: infarmed@infarmed.pt

Design and production: Letras & Sinais, Comunicação e Imagem, Lda.
ISSN: 0873-7118

Alerts and News at the Infarmed website

LinkedIn

Twitter

Facebook

For news and publications,
just use thirty seconds of your time
and register [here!](#)

Risk of Acute Generalized Exanthematous Pustulosis (AGEP) with Acetazolamide and with Macrolides



Quick Read

Acute generalized exanthematous pustulosis is a rare skin adverse reaction that has been associated to various drugs. The anti-glaucoma agent acetazolamide, as well as the macrolide antibiotics are now being added to the list of medicines which can cause this undesirable effect.

Acute generalized exanthematous pustulosis is characterized by non-follicular aseptic pustules covering disseminated skin oedema and erythema. Scaling occurs as the reaction resolves. It usually appears on the face, neck, armpits, groins and other skin folds, and rapidly spreads in a few days to other body areas. The skin manifestations are accompanied by fever higher than 38 degrees Celsius and neutrophilia.

Acetazolamide is a carbonic anhydrase inhibitor with weak diuretic activity. It decreases the production of aqueous humour in the eyes thereby decreasing intraocular pressure. Medicinal products containing acetazolamide are indicated as adjuvants in the treatment of open-angle glaucoma, secondary glaucoma, and closed-angle glaucoma to achieve pre-operative lowering of intraocular pressure.

The macrolides make up a group of broad-spectrum antibiotics with a macrocyclic lactonic ring in their molecular structure, and which inhibit the synthesis of bacterial proteins.

During routine pharmacovigilance activities, the Swedish and the Irish medicines agencies detected a safety signal associated respectively with the use of acetazolamide and macrolides (azithromycin, clarithromycin, erythromycin, roxithromycin) and the seemingly rare occurrence of acute generalized exanthematous pustulosis (AGEP).

Following assessment by the European Medicines Agency (EMA) of EudraVigilance (European ADR report database) cases, literature data concerning both acetazolamide,¹⁻³ and macrolides,⁴⁻¹⁰ as well as complementary data provided by MA holders, changes to SmPCs (and their PILs) will result.

SmPC section 4.8 of all the above products will list AGEP as an undesirable effect.

Section 4.4. on Special warnings and precautions for use will include the following text in the case of ACETAZOLAMIDE:

The occurrence at treatment initiation of a feverish generalised erythema associated with pustula may be a symptom of acute generalised exanthematous pustulosis (AGEP) (see section 4.8). In case of AGEP diagnosis, acetazolamide should be discontinued and any subsequent administration of acetazolamide contraindicated.

In the case of **MACROLIDE ANTIBIOTICS, Section 4.4. on Special warnings and precautions for use** will read slightly different for each individual substance. The text for azithromycin is representative:

[...]

As with erythromycin and other macrolides, rare serious allergic reactions, including angioneurotic oedema and anaphylaxis (rarely fatal), dermatologic reactions including acute generalised exanthematous pustulosis (AGEP), Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) (rarely fatal) and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported. Some of these reactions have resulted in recurrent symptoms and required a longer period of observation and treatment. If an allergic reaction occurs, the drug should be discontinued and appropriate therapy should be instituted. Physicians should be aware that reappearance of the allergic symptoms may occur when symptomatic therapy is discontinued.

References:

- ¹ Paulmann M, Meckenhaupt M. Severe Drug Hypersensitivity Reactions: Clinical Pattern, Diagnosis, Etiology and Therapeutic Options. *Curr Pharm Des.* 2016;22(45) 6852-6861.
- ² Fernando SL. Acute generalized exanthematous pustulosis. *Australas J Dermatol.* 2012 May; 53 (2) 87-92.
- ³ Vassallo C et al. Acute generalized exanthematous pustulosis: report of five cases and systematic review of clinical and histopathological findings. *G Ital Dermatol Venereol.* 2014 Jun; 149 (3): 281-90.
- ⁴ da Cunha Filho RR et al. Acute generalized exanthematous pustulosis by azithromycin. *International Journal of Dermatology* 2015; 54:e233-49.
- ⁵ Yamamoto, H. A Rare Case of a 6-year-old Boy with Acute Generalized Exanthematous Pustulosis. *The Journal of the Japan Pediatric Society.* 2016;120 (2):528.
- ⁶ Fujikura M. A Case of Acute Generalized Exanthematous Pustulosis Preceded by Oral Zithromac. *The 813th Tokyo Regional Meeting of the Japanese Dermatological Association* 2007 Jul 21;12-12.
- ⁷ Zweegers J, Bovenschen HJ. A woman with skin abnormalities around the mouth. *Nederlands Tijdschrift voor Geneeskunde.* 2013;157 (7):311.
- ⁸ Macías J et al. Acute generalized exanthematous pustulosis: A purpose of 4 cases. *Allergy: European Journal of Allergy and Clinical Immunology.* 2016;71(102):417-418.
- ⁹ Sierra Salgado OF et al. Acute generalized exanthematous pustulosis (AGEP) induced by azithromycin. *Allergy: European Journal of Allergy and Clinical Immunology.* 2015; 70 SUPPL. 101:334 on 06 Nov 2015.
- ¹⁰ Kinno M et al. Acute generalized exanthematous pustulosis (AGEP): A case report and literature review. *Journal of Hospital Medicine.* 2012;7:S266.

Communications to Healthcare Professionals published on the Infarmed [website](#)

Click on the links



INN Medicinal product	Target	Communication Online publication date
Bivalirudine Bivalirudina Accord	Physicians: interventional cardiologists.	Dosage regimes approved for percutaneous coronary intervention 11-12-2017
Cladribine Litak Leustatin	Pharmacists: Pharmacists: hospital. Physicians: oncologists and haematologists.	Risk of Progressive Multifocal Leucoencephalopathy (PML) 12-12-2017
Radium 223 dichloride Xofigo	Pharmacists: hospital pharmaceutical service directors. Physicians: oncologists and urologists specialized in prostate cancer; nuclear medicine specialists.	Increased risk of death and fractures in a randomized clinical trial with Xofigo used in association with abiraterone acetate and prednisolone/prednisone 18-12-2017
Bcr-Abl tyrosine kinase inhibitors: Bosutinib, Dasatinib, Imatinib, Nilotinib, Ponatinib Bosulif, Sprycel, Glivec, Tasigna, Iclusig	Pharmacists: hospital pharmaceutical service directors. Physicians: : haematologists and oncologists (in the case of Glivec only).	Need to assess patients for hepatitis B virus before treatment, due to risk of hepatitis B reactivation 04-12-2017
Leuprorelin Eligard	Nurses: hospital and health centres. Pharmacists: : hospital and community. Physicians: urologists, oncologists and internists who may prescribe the product.	Medication errors associated with leakage due to excessive tightening of safety needle 14-12-2017
Saccharomyces boulardii UL-250	Nurses: nursing directors. Pharmacists: hospital pharmaceutical service directors, community pharmacy and OTC product selling point healthcare professionals . Physicians: general/family medicine, internists, infectious diseases specialists, gastroenterologists and oncologists, directors of hospital internal medicine, infectious diseases, gastroenterology, oncology, surgery and intensive care unit dpts.	New contraindication in immunocompromised or seriously ill patients 12-12-2017

Educational Materials published in the Infomed product information [webpage](#)

Click on the links



INN Medicinal product	Target	Communication Online publication date
Apixaban Eliquis	Pharmacists: hospital. Physicians: cardiology, neurology, internal medicine, general/family medicine, haematology/immunotherapy, anaesthesiology, orthopaedics, vascular surgery and gastroenterology.	Prescriber's guide 18-12-2017
Atezolimumab Tecentriq	Physicians: pneumologists experienced in the treatment of lung cancer, urologists experienced in the treatment of bladder cancer, and oncologists experienced in the treatment of lung/bladder cancer.	Guide for healthcare professionals
	Patients	Alert card 05-12-2017
Baricitinib Olumiant	Physicians: rheumatologists and internists.	Prescriber's guide 19-12-2017
Tiotropium bromide Braltus	Physicians: pneumologists, allergy and general/family medicine specialists.	Safety information 19-12-2017
Canacinumab Ilaris	Physicians: rheumatologists and paediatricians.	Guide for administration
	Patients with: Gouty arthritis Still's disease Periodic fever syndromes.	Alert cards: Gouty arthritis Still's disease Periodic fever syndromes 22-12-2017
Cladribine Mavenclad	Physicians: neurologists.	Prescriber's guider
	Patients	Patient's guide 22-12-2017
Sodium colistimethate Colobreathe	Physicians: pneumologists and paediatricians experienced in the treatment of cystic fibrosis.	Physician's guide 19-12-2017
Defibrotide Defitelio	Physicians: prescribers. Other healthcare professionals: transplantation centres.	Patient registry: Letter and Introduction 18-12-2017
Etanercept Enbrel	Patients	Alert card 14-12-2017

Educational Materials published in the Infomed product information webpage

Click on the links



INN Medicinal product	Target	Communication Online publication date
Methylphenidate Concerta, Metilfenidato Mylan, Medikinet, Medicebran, Quasym, Ritalina A, Rubifen	Physicians: paediatricians, neuropaediatricians and psychiatrists.	Prescription guide Checklist before prescribing Checklist for monitoring ongoing therapy Table for monitoring ongoing therapy 22-12-2017
Natalizumab Tysabri	Physicians: : neurologists who treat patients with multiple sclerosis. Patients	Safety information Warning card Forms for beginning continuing discontinuing treatment 22-12-2017
Panobinostat Farydak	Patients	Treatment compliance card 22-12-2017
Ranimizumab Lucentis	Physicians: ophthalmologists Patients	Risk management guide for risks associated with intravitreal injection Treatment guide 12-12-2017
Rivastigmine Rivastigmina Teva	Physicians: neurologists and psychiatrists, general/family medicine, internal medicine.	Memory card for the patient and instructions for use 18-12-2017
Romiplostim Nplate	Physicians: prescribers. Nurses: in charge of reconstituting and administering the product.	Patient selection and training for administration of Nplate® at home 19-12-2017
Voriconazole Voriconazol Aristo	Physicians: haematologists, infectious diseases specialists, dermatologists and STD specialists, oncologists. Patients	Q&A brochure Checklist Alert card 19-12-2017

Compiled by Magda Pedro, José Sequeira

